

Justify all responses with clear explanations and in complete sentences unless otherwise stated. Write up your solutions cleanly and neatly, and clearly identify all problem numbers. Either staple the pages of your assignment together and write your name on the first page, or paperclip the pages and write your name on all pages.

---

**Part I:** No justifications are required for these problems. Answers will be graded on correctness.

1. Identify each of the following statements as true or false:

- (a) Every vector space contains a zero vector.
  - (b) In any vector space,  $\alpha \mathbf{v} = \beta \mathbf{v}$  implies that  $\alpha = \beta$ .
  - (c) In any vector space,  $\alpha \mathbf{v} = \alpha \mathbf{w}$  implies that  $\mathbf{v} = \mathbf{w}$ .
  - (d) If  $U$  is a subspace of  $V$  and  $V$  is a subspace of  $W$ , then  $U$  is a subspace of  $W$ .
  - (e) The empty set is a subspace of any vector space.
  - (f) The intersection of two subspaces is always a subspace.
  - (g) The union of two subspaces is always a subspace.
  - (h) The union of two subspaces is never a subspace.
  - (i) The span of the empty set is the empty set.
  - (j) The span of the zero vector is the zero subspace.
  - (k) If  $S$  is any subset of  $V$ , then  $\text{span}(S)$  is the intersection of all subspaces of  $V$  containing  $S$ .
  - (l) If  $S$  is any subset of  $V$ , then  $\text{span}(S)$  always contains the zero vector.
  - (m) Any set containing the zero vector is linearly independent.
  - (n) Any subset of a linearly independent set is linearly independent.
  - (o) Any subset of a linearly dependent set is linearly dependent.
- 

2. Determine whether or not each given set  $S$  is a subspace of the given vector space  $V$ . For each set that is not a subspace, identify at least one part of the subspace criterion that fails.

- (a)  $V = \mathbb{R}^4$ ,  $S =$  the vectors  $\mathbf{v}$  in  $\mathbb{R}^4$  with  $\mathbf{v} \cdot \langle 1, 0, 1, 1 \rangle = 2$ .
  - (b)  $V = M_{3 \times 3}(\mathbb{R})$ ,  $S =$  the  $3 \times 3$  matrices with nonnegative real entries.
  - (c)  $V =$  real-valued functions on  $[0, 1]$ ,  $S =$  the functions with  $f''(x) = f(x)$ .
  - (d)  $V = \mathbb{C}^5$ ,  $S =$  the vectors  $\langle a, b, c, d, e \rangle$  with  $e = a + b$  and  $b = c = d$ .
  - (e)  $V =$  real-valued functions on  $\mathbb{R}$ ,  $S =$  the functions with  $f(x) = f(1 - x)$  for all real  $x$ .
  - (f)  $V =$  real-valued functions on  $\mathbb{R}$ ,  $S =$  the functions that are strictly increasing.
  - (g)  $V = P_3(\mathbb{C})$ ,  $S =$  the polynomials in  $V$  with  $p(i) = 0$ .
  - (h)  $V = M_{2 \times 2}(\mathbb{Q})$ ,  $S =$  the matrices in  $V$  of determinant zero.
- 

3. For each set of vectors in each vector space, determine if they (i) span  $V$ , (ii) are linearly independent, and (iii) are a basis:

- (a)  $\langle 1, 2 \rangle, \langle 3, 2 \rangle, \langle 1, 1 \rangle$  in  $\mathbb{R}^2$ .
  - (b)  $\langle 1, 2, 4 \rangle, \langle 3, 2, 1 \rangle, \langle 1, 1, 1 \rangle$  in  $\mathbb{R}^3$ .
  - (c)  $1 + x, x + x^2$  in  $P_2(\mathbb{C})$ .
  - (d)  $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}$  in  $M_{2 \times 2}(\mathbb{F}_5)$ .
  - (e)  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  in  $M_{2 \times 2}(\mathbb{F}_2)$ .
-

**Part II:** Solve the following problems. Justify all answers with rigorous, clear explanations.

4. Suppose  $A$  is an  $m \times n$  matrix with entries from the field  $F$ .

- (a) Show that the set of all vectors  $\mathbf{x} \in F^m$  such that  $A\mathbf{x}$  equals the zero vector (in  $F^m$ ) is a subspace of  $F^m$ .
  - (b) Deduce that the set of solutions to any homogeneous system of linear equations (i.e., in which all of the constants are equal to zero) over  $F$  is an  $F$ -vector space.
- 

5. Suppose  $V$  is a vector space and let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  and  $T = \{\mathbf{v}_1, \mathbf{v}_1 + \mathbf{v}_2, \mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3\}$ .

- (a) If  $S$  is linearly independent, show that  $T$  is linearly independent.
  - (b) If  $S$  spans  $V$ , show that  $T$  spans  $V$ .
  - (c) If  $S$  is a basis for  $V$ , show that  $T$  is a basis for  $V$ .
- 

6. If  $V$  is a vector space and  $W_1, W_2$  are two subspaces of  $V$ , their sum is defined to be the set  $W_1 + W_2 = \{\mathbf{w}_1 + \mathbf{w}_2 : \mathbf{w}_1 \in W_1 \text{ and } \mathbf{w}_2 \in W_2\}$  of all sums of an element of  $W_1$  with an element of  $W_2$ .

- (a) Prove that  $W_1 + W_2$  contains  $W_1$  and  $W_2$ , and is a subspace of  $V$ .
  - (b) Prove in fact that  $W_1 + W_2$  is the smallest subspace containing both  $W_1$  and  $W_2$ .
  - (c) Show that if  $S_1$  and  $S_2$  are arbitrary subsets of  $V$ , then  $\text{span}(S_1 \cup S_2) = \text{span}(S_1) + \text{span}(S_2)$ .
  - (d) For  $V = F[x]$ , let  $W_1$  be the subspace of all even polynomials (i.e., polynomials with all terms of even degree) and  $W_2$  be the subspace of all odd polynomials (polynomials with all terms of odd degree). Show that  $V = W_1 + W_2$ .
- 

7. Suppose that  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  and  $\{\mathbf{w}_1, \mathbf{w}_2\}$  are two linearly-independent subsets of the vector space  $V$ .

- (a) If  $\text{span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) \cap \text{span}(\mathbf{w}_1, \mathbf{w}_2) = \{\mathbf{0}\}$ , prove that  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}_1, \mathbf{w}_2\}$  is a linearly independent set.
  - (b) If  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}_1, \mathbf{w}_2\}$  is a linearly independent set, prove that  $\text{span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) \cap \text{span}(\mathbf{w}_1, \mathbf{w}_2) = \{\mathbf{0}\}$ .
-