

Question:	1	2	3	4	5	6	7	8	9	10	11	Total
Points:	6	8	10	10	8	10	8	10	10	10	10	100
Score:												

**PRINT** Your name here: \_\_\_\_\_

**PRINT** Your instructor's name here: \_\_\_\_\_

**Please check that you have 11 different pages.**

**Answers from your calculator, without supporting work, are worth zero points.**

1. (6 points) Find an equation for the tangent plane to the graph of  $f(x, y) = e^{2(y-1)}\sqrt{x}$  at the point  $(x, y) = (4, 1)$ .

2. An electric dipole is located at the origin in the  $xy$ -plane and generates an electrostatic potential given by

$$V(x, y) = \frac{y}{x^2 + y^2}.$$

Here  $V$  is measured in volts and  $x$  and  $y$  are measured in meters.

- (a) (4 points) What is the gradient of the potential  $V$  at the point  $(x, y) = (1, 2)$ ?

- (b) (4 points) The level curves of  $V$  are called equipotential curves. At the point  $(1, 2)$ , find the direction of a vector which is tangent to the equipotential curve passing through that point. Give your answer as a unit vector with positive  $\vec{i}$ -component.

3. (10 points) Find the critical points of  $f(x, y) = x^3 - 3xy + y^3$ , and classify each critical point as a point where  $f$  has a local maximum value, a local minimum value, or a saddle point.

4. (10 points) A box-shaped building with a rectangular base is to have a volume of 8000  $\text{ft}^3$ . Suppose that annual heating and cooling costs will amount to  $\$2/\text{ft}^2$  for its roof, front wall, and back wall, and  $\$4/\text{ft}^2$  for the two remaining walls. Note that the floor is excluded. What dimensions of the building would minimize these annual costs?

5. (8 points) Let  $R$  be the filled-in triangle in the first quadrant of the  $xy$ -plane which is bounded by the  $y$ -axis, the line where  $y = x$ , and the line where  $y = 1$ . Let  $T$  be the solid region in  $\mathbb{R}^3$  which lies above  $R$  and below the surface where  $z = 12y^2 - 12x^2 + 24$ . Find the volume of the solid region  $T$ .

6. (10 points) Evaluate the integral

$$\int_0^{\frac{3}{\sqrt{2}}} \int_x^{\sqrt{9-x^2}} \int_0^{9-x^2-y^2} y \, dz \, dy \, dx.$$

7. (8 points) Let  $S$  be the portion of the cone given by  $\phi = 5\pi/6$  that is inside the sphere of radius 5. This surface can be parameterized by

$$\vec{r}(u, v) = \left( \frac{1}{2}u \cos v, \frac{1}{2}u \sin v, -\frac{\sqrt{3}}{2}u \right), \quad 0 \leq u \leq 5, \quad 0 \leq v \leq 2\pi.$$

Find the surface area of  $S$ .


8. Consider the vector field  $\mathbf{F}(x, y) = (2x^3y^4 + x, 2x^4y^3 + y^2)$  in  $\mathbb{R}^2$ .
- (a) (2 points) Show that  $\mathbf{F}$  is conservative.
- (b) (4 points) Find a potential function  $f$  of  $\mathbf{F}$ . Show your work.
- (c) (4 points) Compute the integral  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where  $\mathbf{F}$  is the vector field from part (a) and  $C$  is the half-circle of radius 2 centered at the origin and going clockwise from  $(0, -2)$  to  $(0, 2)$ .



9. (10 points) Consider the vector field given by  $\mathbf{F}(x, y) = (\sqrt{1+x^3}, 2xy)$ . Calculate

$$\int_C \mathbf{F} \cdot d\mathbf{r}$$

where  $C$  is the curve which starts at  $(0, 0)$ , then goes along the  $x$ -axis until it reaches  $(2, 0)$ , then goes vertically until it reaches  $(2, 1)$ , and then goes horizontally until it reaches  $(0, 1)$ .



10. (10 points) Use the Divergence theorem to find the flux of the vector field

$$\mathbf{F}(x, y, z) = (0, 0, z^3/3)$$

across the sphere of radius 1 centered at the origin oriented by the outward pointing normal.

11. Let  $M$  be the portion of the graph of  $z = 1 - x^2 - y^2$  that lies above the  $xy$ -plane ( $z \geq 0$ ), oriented upward. Let

$$\mathbf{F}(x, y, z) = (x^2 + y^2 + y + z^2, 3z, 5 - x^2 - y^2).$$

- (a) (3 points) Let  $\partial M$  be the boundary of  $M$ . Write a parameterization of  $\partial M$ .

- (b) (7 points) Use Stokes' theorem to compute

$$\iint_M (\vec{\nabla} \times \mathbf{F}) \cdot \vec{n} \, dS.$$

(Hint: Depending on how you solve the problem, the following fact may (or may not) be useful:  $\int_0^{2\pi} (\sin x)^2 \, dx = \pi$ .) Be careful to explain each step in your calculation.