

Justify all responses with clear explanations and in complete sentences unless otherwise stated. Write up your solutions cleanly and neatly, and clearly identify all problem numbers. Identify all pages containing each problem when submitting the assignment.

Part I: No justifications are required for these problems. Answers will be graded on correctness.

1. In each given quadratic integer ring, determine which of the given elements are units, which are irreducible, and which are reducible. Also, for the units, compute their multiplicative inverses, and for the reducible elements find a nontrivial factorization.

(a) $R = \mathbb{Z}[i]$, elements $4 - i, 3 + i, 3 - 2i, 7$.

(b) $R = \mathcal{O}_{\sqrt{-3}}$, elements $\frac{1 + \sqrt{-3}}{2}, 2 + \sqrt{-3}, 3 + \sqrt{-3}, \frac{5 + \sqrt{-3}}{2}$.

(c) $R = \mathcal{O}_{\sqrt{5}}$, elements $2 + \sqrt{5}, 3 - 2\sqrt{5}, 7 + 5\sqrt{5}, 1 + \sqrt{5}$.

(d) $R = \mathcal{O}_{\sqrt{7}}$, elements $2 - \sqrt{7}, 3 + \sqrt{7}, 1 + \sqrt{7}, 8 - 3\sqrt{7}$.

2. For each pair of elements a, b in the given Euclidean domain R , find a greatest common divisor d and write it in the form $d = xa + yb$ for some $x, y \in R$. (You may wish to work through problems 4 and 5 before doing parts (c), (d), and (e).)

(a) $R = \mathbb{Z}[i]$, $a = 57 + 17i$, $b = 26 + 22i$.

(b) $R = \mathbb{Z}[i]$, $a = 9 + 43i$, $b = 22 + 10i$.

(c) $R = \mathbb{Z}[\sqrt{-2}]$, $a = 33 + 5\sqrt{-2}$, $b = 8 + 11\sqrt{-2}$.

(d) $R = \mathbb{Z}[\sqrt{2}]$, $a = 31 + 15\sqrt{2}$, $b = 10 + \sqrt{2}$.

(e) $R = \mathcal{O}_{\sqrt{-3}}$, $a = 19 + \sqrt{-3}$, $b = 14 + 7\sqrt{-3}$.

Part II: Solve the following problems. Justify all answers with rigorous, clear arguments.

3. Show that the rings $(\mathbb{Z}/15\mathbb{Z}) \times (\mathbb{Z}/8\mathbb{Z})$ and $(\mathbb{Z}/24\mathbb{Z}) \times (\mathbb{Z}/5\mathbb{Z})$ are isomorphic.
-

4. Let $R = \mathbb{Z}[\sqrt{-2}]$, and let $a + b\sqrt{-2}$ and $c + d\sqrt{-2}$ be elements of R with $c + d\sqrt{-2} \neq 0$.

(a) Show that $\frac{a + b\sqrt{-2}}{c + d\sqrt{-2}} = x + y\sqrt{-2}$ for rational x, y . Then let s be the closest integer to x and t be the closest integer to y , and set $q = s + t\sqrt{-2}$ and $r = (a + b\sqrt{-2}) - (s + t\sqrt{-2})(c + d\sqrt{-2})$. Prove also that $N(r) \leq \frac{3}{4}N(c + d\sqrt{-2})$.

(b) Show that R is a Euclidean domain.

(c) Show that $\mathbb{Z}[\sqrt{2}]$ and $\mathbb{Z}[\sqrt{3}]$ are also Euclidean domains under the absolute value of the field norm $|N(a + b\sqrt{D})| = |a^2 - Db^2|$.

5. The goal of this problem is to prove that $\mathcal{O}_{\sqrt{-D}}$ is a Euclidean domain for $-D = -3, -7,$ and $-11,$ which extends the result of problem 4 (establishing this fact for $-D = -2, 2,$ and 3).

- (a) Suppose ABC is a triangle. Show that the point P inside ABC that maximizes the distance to the nearest vertex of ABC is the circumcenter (i.e., the center of the circle through the vertices of ABC , or equivalently, the point O such that $OA = OB = OC$).
 - (b) Suppose that $-D = -3, -7,$ or $-11.$ Prove that any complex number $z \in \mathbb{C}$ differs from an element in $\mathcal{O}_{\sqrt{-D}}$ by a complex number whose norm (i.e., the square of its absolute value) is at most $\frac{(1+D)^2}{16D}.$ [Hint: The elements of $\mathcal{O}_{\sqrt{-D}}$ form a lattice Λ in the complex plane. Identify a fundamental region for this lattice and then use symmetry to reduce the minimal distance calculation to part (a).]
 - (c) Prove that $\mathcal{O}_{\sqrt{-D}}$ is a Euclidean domain for $-D = -3, -7,$ and $-11.$ [Hint: Adapt the proof in 4b.]
-

6. The goal of this problem is to prove that for any squarefree integer $D \geq 3,$ the ring $\mathbb{Z}[\sqrt{-D}]$ is not a unique factorization domain, generalizing the technique used for $D = 5.$

- (a) Show that $\sqrt{-D}, 1 + \sqrt{-D}, 1 - \sqrt{-D},$ and 2 are irreducible elements in $\mathbb{Z}[\sqrt{-D}].$ [Hint: For the first three, show that the only elements of norm less than D are integers.]
 - (b) Show that either D (if D is even) or $D + 1$ (if D is odd) has two different factorizations into irreducibles in $\mathbb{Z}[\sqrt{-D}],$ and deduce that $\mathbb{Z}[\sqrt{-D}]$ is not a unique factorization domain.
 - (c) What goes wrong if you try to use the proof to show that $\mathbb{Z}[\sqrt{D}]$ is not a UFD for squarefree $D \geq 3?$
-

7. [Challenge] Let $R = \mathbb{Z}[\sqrt{-3}]$ and let $I = (2, 1 + \sqrt{-3})$ in $R.$

- (a) Show that $I^2 = (2)I$ in R but that $I \neq (2).$
- (b) Show that there are two residue classes in R/I and deduce that I is a prime ideal.
- (c) Show that I is the unique proper ideal of R properly containing (2) and also the unique prime ideal of R containing $(2).$ [Hint: Consider the ideals of $R/(2)$ and use the correspondence between ideals of R containing J and ideals of $R/J.$]
- (d) Show that (2) cannot be written as a product of prime ideals of $R.$

Remark: This problem illustrates that factorization into prime ideals can fail if we do not work in the full quadratic integer ring. Working in the correct ring $\mathcal{O}_{\sqrt{-3}} = \mathbb{Z}[\frac{1+\sqrt{-3}}{2}]$ will solve the issues that arise in this example, since in fact $I = (2)$ is a prime ideal inside $\mathcal{O}_{\sqrt{-3}}$ because 2 and $1 + \sqrt{-3}$ are now associates.
