

1. For vectors  $\mathbf{v} = \langle 3, 0, -4 \rangle$  and  $\mathbf{w} = \langle -1, 6, 2 \rangle$ , compute the following:
- (a)  $\mathbf{v} + 2\mathbf{w}$ ,  $\|\mathbf{v}\|$ , and  $\|\mathbf{w}\|$ .
  - (b)  $\mathbf{v} \cdot \mathbf{w}$  and  $\mathbf{v} \times \mathbf{w}$ .
  - (c) A unit vector in the opposite direction of  $\mathbf{v}$ .
  - (d) A vector of length 4 in the same direction as  $\mathbf{w}$ .
  - (e) The angle between  $\mathbf{v}$  and  $\mathbf{w}$ .
  - (f) The area of the parallelogram with sides  $\mathbf{v}$ ,  $\mathbf{w}$ .
- 

2. Find an equation or a parametrization for each of the following:
- (a) The plane parallel to  $x + 2y - 3z = 1$  containing the point  $(2, -1, 2)$ .
  - (b) The line containing  $(2, -1, 4)$  and  $(3, 6, 2)$ .
  - (c) The line parallel to  $\langle x, y, z \rangle = \langle 1 - 2t, 3 + 2t, 2 + 5t \rangle$  containing the point  $(1, 1, 1)$ .
  - (d) The plane perpendicular to the line  $\langle x, y, z \rangle = \langle t, 1 + 2t, 3 - 3t \rangle$  containing the origin.
  - (e) The plane containing the points  $(1, 0, 1)$ ,  $(2, 1, 2)$ , and  $(3, 3, 5)$ .
  - (f) The intersection of the planes  $x + y + 2z = 4$  and  $2x - y - z = 5$ .
  - (g) The plane containing the vectors  $\langle 1, 2, -1 \rangle$  and  $\langle 2, -1, 1 \rangle$  and the point  $(1, -1, 2)$ .
- 

3. At time  $t$ , a particle has position  $\mathbf{r}(t) = \langle 3 \cos(t), 5 \sin(t), 4 \cos(t) \rangle$ . Compute the following:
- (a) The particle's velocity  $\mathbf{v}(t)$ .
  - (b) The particle's speed  $\|\mathbf{v}(t)\|$ .
  - (c) The distance traveled by the particle between  $t = 0$  and  $t = 1$ .
  - (d) The particle's acceleration  $\mathbf{a}(t)$ .
  - (e) The unit tangent vector  $\mathbf{T}(t)$ .
  - (f) The unit normal vector  $\mathbf{N}(t)$ .
- 

4. Find a parametrization for the tangent line to the curve  $\langle x, y, z \rangle = s^2\mathbf{i} + s^3\mathbf{j} + s^4\mathbf{k}$  when  $s = 1$ .
- 

5. A potato is fired into the air at time  $t = 0$ s from the origin in a vacuum with initial velocity  $\mathbf{v}(0) = (4\mathbf{i} + 8\mathbf{j} + 80\mathbf{k})\text{m/s}$ . Assuming that the only force acting on the potato is the downward acceleration due to gravity of  $\mathbf{a}(t) = -10\mathbf{k}\text{m/s}^2$ , find the following:
- (a) The velocity of the potato at time  $t$ .
  - (b) The position of the potato at time  $t$ .
  - (c) The total time that the potato is in the air.
  - (d) The potato's speed when it hits the ground.
- 

6. If  $f(x, y) = 3x^2e^{xy}$ , find  $\frac{\partial f}{\partial x}$ ,  $f_y$ ,  $f_{xx}$ ,  $\frac{\partial^2}{\partial y \partial x} f$ ,  $f_{yy}$ , and  $f_{yyyy}(1, 2)$ .
-

7. Let  $f(x, y, z) = x^3yz^2$  and  $g(x, y, z) = \ln(x^2 + y^2 + z^2)$ .
- (a) Find the rates of change of  $f$  and  $g$  in the direction of  $\mathbf{v} = \langle 2, -1, 2 \rangle$  at the point  $(1, 1, 1)$ .
  - (b) Find the minimum and maximum rates of change of  $f$  and  $g$  at the point  $(1, 2, 1)$ , and the unit vector directions in which the minimum and maximum rates of change occur.
  - (c) Find the linearizations to  $f$  and  $g$  at the point  $(2, 1, 1)$ .
- 

8. Consider the surface  $e^{x-yz} + xz = 9$ .

- (a) Find an equation for the tangent plane to the surface at  $(4, 2, 2)$ .
  - (b) Assuming that  $z$  is defined implicitly as a function of  $x$  and  $y$ , find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$ .
- 

9. Suppose  $f(x, y)$  has  $\frac{\partial f}{\partial x}(1, 5) = 9$ ,  $\frac{\partial f}{\partial y}(1, 5) = -3$ ,  $\frac{\partial f}{\partial x}(2, -2) = 4$ , and  $\frac{\partial f}{\partial y}(2, -2) = 5$ , where  $x(s, t)$  and  $y(s, t)$  are such that  $x(1, 5) = 2$ ,  $y(1, 5) = -2$ ,  $\frac{\partial x}{\partial s}(1, 5) = 3$ ,  $\frac{\partial x}{\partial t}(1, 5) = 2$ ,  $\frac{\partial y}{\partial s}(1, 5) = 4$ , and  $\frac{\partial y}{\partial t}(1, 5) = -2$ . Find:

- (a)  $\frac{\partial f}{\partial s}$  at  $(s, t) = (1, 5)$ .
  - (b)  $\frac{\partial f}{\partial t}$  at  $(s, t) = (1, 5)$ .
- 

10. Suppose  $f(x, y) = x^3 + 3xy$ .

- (a) Find  $f_{xxy}$ .
  - (b) Find the rate of change of  $f$  at the point  $(x, y) = (1, 2)$  in the direction of the origin.
  - (c) Find the direction in which  $f$  is decreasing the fastest at  $(x, y) = (2, 0)$ .
  - (d) If  $x = 3s^2 - 2t$  and  $y = s^3t$ , find  $\partial f / \partial t$  when  $s = 1$  and  $t = 2$ .
  - (e) Find the linearization of  $f$  at  $(x, y) = (1, 3)$  and use it to estimate  $f(1.01, 3.02)$ .
  - (f) Find an equation for the tangent plane to  $z = f(x, y)$  at  $(x, y) = (-1, 1)$ .
  - (g) Find and classify the critical points for  $f$ .
- 

11. Find and classify the critical points for each function as minima, maxima, or saddle points:

- (a)  $f(x, y) = xy - x^2 - y^2 - 2x - 2y$ .
  - (b)  $f(x, y) = x^4 + y^2 - 8x^2 + 4y$ .
  - (c)  $f(x, y) = xy + \frac{1}{x} + \frac{1}{y}$ .
  - (d)  $f(x, y) = x^3 - 3xy + 3y^2$ .
-