

1. (a) Take $\mathbf{r}(t) = \langle 3t, 1+t \rangle$ for $0 \leq t \leq 1$ with $ds = \|\mathbf{v}(t)\| dt = \sqrt{10} dt$. Integral is $\int_0^1 (3t)^2 \sqrt{10} dt = [3\sqrt{10}]$.
- (b) Take $\mathbf{r}(t) = \langle 3 \cos t, 3 \sin t \rangle$ for $0 \leq t \leq 2\pi$ with $dx = -3 \sin t dt$ and $dy = 3 \cos t dt$. Integral is $\int_0^{2\pi} (3 \cos t)(-3 \sin t) dt + (3 \sin t)(3 \cos t) dt = \int_0^{2\pi} 0 dt = [0]$.
- (c) Have $ds = \|\mathbf{v}(t)\| = \|\langle -3 \sin 3t, 3 \cos 3t, 4 \rangle\| = 5$ and $\sqrt{x^2 + y^2} = 1$. Integral is $\int_0^\pi 1 \cdot 5 dt = [5\pi]$.
- (d) Average value is $[\int_C y ds] / [\int_C 1 ds]$. Parametrize by $\mathbf{r}(t) = \langle 2 \cos t, 2 \sin t \rangle$ for $0 \leq t \leq \pi$ with $ds = \|\mathbf{v}(t)\| dt = 2 dt$. Numerator is $\int_0^\pi 2 \sin t \cdot 2 dt = 8$, denominator is $\int_0^\pi 2 dt = 2\pi$, ratio is $[8/(2\pi) = 4/\pi]$.
- (e) Work is $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P dx + Q dy + R dz$. With $x = t^2$, $y = t^4$, $z = t^3$ get $dx = 2t dt$, $dy = 4t^3 dt$, $dz = 3t^2 dt$, $P = yz = t^7$, $Q = xz = t^5$, $R = x^3 = t^6$. Integral is $\int_0^1 t^7 \cdot 2t dt + t^5 \cdot 4t^3 dt + t^6 \cdot 3t^2 dt = \int_0^1 9t^8 dt = [1 \text{ J}]$.
- (f) Parametrize by $\mathbf{r}(t) = \langle \cos t, \sin t \rangle$ for $0 \leq t \leq \pi$. With $x = \cos t$, $y = \sin t$, get $dx = -\sin t dt$, $dy = \cos t dt$, $P = y + x = \sin t + \cos t$, $Q = y - x = \sin t - \cos t$. Circulation is $\int_C P dx + Q dy = \int_0^\pi (\sin t + \cos t)(-\sin t dt) + (\sin t - \cos t)(\cos t dt) = \int_0^\pi -1 dt = [-\pi]$. Flux is $\int_C -Q dx + P dy = \int_0^\pi (\cos t - \sin t)(-\sin t dt) + (\sin t + \cos t)(\cos t dt) = \int_0^\pi 1 dt = [\pi]$.
- (g) With $x = t^2$, $y = t^3$ get $dx = 2t dt$, $dy = 3t^2 dt$, $P = xy = t^5$, $Q = y^2 = t^6$. Circulation is $\int_C P dx + Q dy = \int_0^1 t^5(2t dt) + t^6(3t^2 dt) = \int_0^1 (2t^6 + 3t^8) dt = [13/21]$. Flux is $\int_C -Q dx + P dy = \int_0^1 -t^6(2t dt) + t^5(3t^2 dt) = \int_0^1 t^7 dt = [1/8]$.
- (h) Have $ds = \|\mathbf{v}(t)\| = \langle -\pi \sin \pi t, \pi \cos \pi t, 2 \rangle = \sqrt{\pi^2 + 4}$ and $x^2 + y^2 + z^3 = 1 + 8t^3$. Integral is $\int_0^2 (1 + 8t^3) \sqrt{\pi^2 + 4} dt = [34\sqrt{\pi^2 + 4}]$.
- (i) Work is $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P dx + Q dy + R dz$. Get $dx = \cos t dt$, $dy = 3dt$, $dz = -3 \sin t dt$ so $P = 5$, $Q = \sin^2 t$, $R = -3 \cos t$. Integral is $\int_0^{2\pi} 5 \cdot \cos t dt + (\cos^2 t) \cdot 3 dt - 3 \sin t \cdot (-3 \sin t dt) = \int_0^{2\pi} (5 \cos t + 9) dt = [18\pi]$.
-
2. (a) Writing as $(x/2)^2 + y^2 = 1$ we can parametrize as $x/2 = \cos t$, $y = \sin t$ for $0 \leq t \leq \pi$, which gives $\mathbf{r}(t) = \langle 2 \cos t, \sin t \rangle$ for $0 \leq t \leq \pi$.
- (b) Circulation is $\int_C P dx + Q dy = \int_0^{2\pi} [2 \sin^2 t + 2 \cos^2 t + 2 \sin t \cos t] dt = \int_0^{2\pi} [2 + \sin 2t] dt = [4\pi]$.
- (c) Flux is $\int_C -Q dx + P dy = \int_0^{2\pi} [4 \sin^2 t \cos t] dt = \frac{4}{3} \sin^3 t \Big|_{t=0}^{2\pi} = [0]$.
-
3. (a) Normal vector is $\mathbf{n} = (\mathbf{r}/ds) \times (\mathbf{r}/dt) = \langle 1, 2s, 3s^2 \rangle \times \langle 1, 2t, 3t^2 \rangle$. With $s = 1$ and $t = 2$ this is $\mathbf{n} = \langle 1, 2, 3 \rangle \times \langle 1, 4, 12 \rangle = \langle 12, -9, 2 \rangle$. Point is $\mathbf{r}(1, 2) = \langle 3, 5, 9 \rangle$, so equation is $[12(x - 3) - 9(y - 5) + 2(z - 9) = 0]$.
- (b) If $(x, y, z) = (1, 2, -3)$ then $t = -1$ and $s = -1$. Then $\mathbf{n} = (\mathbf{r}/ds) \times (\mathbf{r}/dt) = \langle 2s, 2t, 0 \rangle \times \langle 0, 2s, 9t^2 \rangle$, so $s = -1$ and $t = -1$ give $\mathbf{n} = \langle -2, -2, 0 \rangle \times \langle 0, -2, 9 \rangle = \langle -18, 18, 4 \rangle$. Equation is $[-18(x - 1) + 18(y - 2) + 4(z + 3) = 0]$.
-
4. (a) Parametrize by $\mathbf{r}(s, t) = \langle s, t, 2s + 2t \rangle$ for $1 \leq s \leq 2$, $2 \leq t \leq 4$. Then $\frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} = \langle -2, -2, 1 \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} \right\| ds dt = 3 ds dt$. Surface integral is $\int_2^4 \int_1^2 (s^2 + t^2) \cdot 3 ds dt$.
- (b) Parametrize by $\mathbf{r}(z, \theta) = \langle 2 \cos \theta, 2 \sin \theta, z \rangle$ for $0 \leq \theta \leq 2\pi$, $0 \leq z \leq 4$. Then $\frac{\partial \mathbf{r}}{\partial z} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle -2 \cos \theta, 2 \sin \theta, 0 \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial z} \times \frac{\partial \mathbf{r}}{\partial \theta} \right\| dz d\theta = 2 dz d\theta$. Surface integral is $\int_0^{2\pi} \int_0^4 8z dz d\theta$.
- (c) Parametrize by $\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, 4r \rangle$ for $0 \leq \theta \leq 2\pi$, $0 \leq r \leq 2$. Then $\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle -4r \cos \theta, -4r \sin \theta, r \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} \right\| dr d\theta = r \sqrt{17} dr d\theta$. Surface integral is $\int_0^{2\pi} \int_0^2 r \cdot r \sqrt{17} dr d\theta$.
- (d) Parametrize by $\mathbf{r}(\theta, \varphi) = \langle \cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi \rangle$ for $0 \leq \theta \leq 2\pi$, $0 \leq \varphi \leq \pi$. Then $\frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \varphi} = \langle -\cos \theta \sin^2 \varphi, -\sin \theta \sin^2 \varphi, -\sin \varphi \cos \varphi \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \varphi} \right\| d\varphi d\theta = \sin \varphi d\varphi d\theta$. Surface integral is $\int_0^{2\pi} \int_0^\pi \cos^2 \varphi \sin \varphi d\varphi d\theta$.
- (e) Parametrize by $\mathbf{r}(s, t) = \langle s, t, s^2 + t^2 \rangle$ for $0 \leq s \leq 1$, $0 \leq t \leq 1$. Then $\frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} = \langle -2s, -2t, 1 \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} \right\| ds dt = \sqrt{4s^2 + 4t^2 + 1} ds dt$. Surface integral is $\int_0^1 \int_0^1 s \sqrt{4s^2 + 4t^2 + 1} ds dt$.
- (f) Parametrize by $\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, r^2 \rangle$ for $0 \leq \theta \leq 2\pi$, $0 \leq r \leq 3$. Then $\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle -2r^2 \cos \theta, -2r^2 \sin \theta, r \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} \right\| dr d\theta = r \sqrt{4r^2 + 1} dr d\theta$. Surface integral is $\int_0^{2\pi} \int_0^3 r \cdot r \sqrt{4r^2 + 1} dr d\theta$.

- (g) Parametrize by $\mathbf{r}(\theta, \varphi) = \langle 3 \cos \theta \sin \varphi, 3 \sin \theta \sin \varphi, 3 \cos \varphi \rangle$ for $0 \leq \theta \leq \pi/2, 0 \leq \varphi \leq \pi/2$. Then $\frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \varphi} = \langle -9 \cos \theta \sin^2 \varphi, -9 \sin \theta \sin^2 \varphi, -9 \sin \varphi \cos \varphi \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \varphi} \right\| d\varphi d\theta = 9 \sin \varphi d\varphi d\theta$. Surface integral is $\int_0^{\pi/2} \int_0^{\pi/2} 3 \sin \theta \sin \varphi \cdot 9 \sin \varphi d\varphi d\theta$.
-
5. (a) Parametrize by $\mathbf{r}(s, t) = \langle s, t, 3s + 4t + 11 \rangle$ for $0 \leq s \leq 1, 0 \leq t \leq 2$. Then $\frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} = \langle -3, -4, 1 \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} \right\| ds dt = \sqrt{26} ds dt$. Surface area integral is $\int_0^1 \int_0^2 1 \cdot \sqrt{26} ds dt$.
- (b) Here $\frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} = \langle 2s, t, 0 \rangle \times \langle 0, s, 2t \rangle = \langle 2t^2, -4st, 2s^2 \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} \right\| ds dt = 2\sqrt{s^4 + 4s^2t^2 + t^4} ds dt$. Surface area integral is $\int_0^1 \int_0^2 1 \cdot 2\sqrt{s^4 + 4s^2t^2 + t^4} ds dt$.
- (c) Parametrize by $\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, 3r \rangle$ for $0 \leq \theta \leq 2\pi, 2 \leq r \leq 3$. Then $\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle -3r \cos \theta, 3r \sin \theta, r \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} \right\| dr d\theta = r\sqrt{10} dr d\theta$. Surface area integral is $\int_0^{2\pi} \int_2^3 1 \cdot r\sqrt{10} dr d\theta$.
- (d) Cone is $\varphi = \pi/6$, so can parametrize by $\mathbf{r}(\theta, \varphi) = \langle 3 \cos \theta \sin \varphi, 3 \sin \theta \sin \varphi, 3 \cos \varphi \rangle$ for $0 \leq \theta \leq 2\pi, \pi/6 \leq \varphi \leq \pi$. Then $\frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \varphi} = \langle -9 \cos \theta \sin^2 \varphi, -9 \sin \theta \sin^2 \varphi, -9 \sin \varphi \cos \varphi \rangle$ so $d\sigma = \left\| \frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \varphi} \right\| d\varphi d\theta = 9 \sin \varphi d\varphi d\theta$. Surface area integral is $\int_0^{2\pi} \int_{\pi/6}^{\pi} 1 \cdot 9 \sin \varphi d\varphi d\theta$.
-
6. (a) Parametrize as $\mathbf{r}(s, t) = \langle s, t, 4 - s - t \rangle$ for $0 \leq s \leq 1, 0 \leq t \leq 2$. Then $\frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} = \langle 1, 1, 1 \rangle$ and $\mathbf{F} = \langle s^2, 2s^2, 3s^2 \rangle$. Surface integral is $\int_0^2 \int_0^1 \langle 1, 1, 1 \rangle \cdot \langle s^2, 2s^2, 3s^2 \rangle ds dt = \int_0^2 \int_0^1 6s^2 ds dt = \boxed{4}$.
- (b) Parametrize as $\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, r \rangle$ for $0 \leq r \leq \sqrt{6}, 0 \leq \theta \leq 2\pi$. Then $\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle -r \cos \theta, -r \sin \theta, r \rangle$ and $\mathbf{F} = \langle -r^2 \cos \theta, -r^2 \sin \theta, r^2 \rangle$. Surface integral is $\int_0^{2\pi} \int_0^{\sqrt{6}} \langle -r \cos \theta, -r \sin \theta, r \rangle \cdot \langle -r^2 \cos \theta, -r^2 \sin \theta, r^2 \rangle dr d\theta = \int_0^{2\pi} \int_0^{\sqrt{6}} 2r^3 dr d\theta = \boxed{36\pi}$.
- (c) Parametrize as $\mathbf{r}(\theta, z) = \langle 2 \cos \theta, 2 \sin \theta, z \rangle$ for $0 \leq \theta \leq 2\pi, 0 \leq z \leq 3$. Then $\frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial z} = \langle 2 \cos \theta, 2 \sin \theta, 0 \rangle$ and $\mathbf{F} = \langle 2z \cos \theta, 2z \sin \theta, z^4 \rangle$. Surface integral is $\int_0^{2\pi} \int_0^3 \langle 2 \cos \theta, 2 \sin \theta, 0 \rangle \cdot \langle 2z \cos \theta, 2z \sin \theta, z^4 \rangle dz d\theta = \int_0^{2\pi} \int_0^3 4z dz d\theta = \boxed{36\pi}$.
- (d) Parametrize as $\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, r^2 \rangle$ for $0 \leq \theta \leq 2\pi, 0 \leq r \leq \sqrt{2}$. Then $\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle -2r^2 \cos \theta, -2r^2 \sin \theta, r \rangle$ and $\mathbf{F} = \langle r \sin \theta, -r \cos \theta, r^2 \rangle$. Surface integral is $\int_0^{2\pi} \int_0^{\sqrt{2}} \langle -2r^2 \cos \theta, -2r^2 \sin \theta, r \rangle \cdot \langle r \sin \theta, -r \cos \theta, r^2 \rangle dr d\theta = \int_0^{2\pi} \int_0^{\sqrt{2}} r^3 dr d\theta = \boxed{2\pi}$.
- (e) Parametrize by $\mathbf{r}(\theta, \varphi) = \langle \cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi \rangle$ for $0 \leq \theta \leq 2\pi, 0 \leq \varphi \leq \pi/2$. Then $\frac{\partial \mathbf{r}}{\partial \varphi} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle \cos \theta \sin^2 \varphi, \sin \theta \sin^2 \varphi, \sin \varphi \cos \varphi \rangle$ and $\mathbf{F} = \langle \cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi \rangle$. Integral is then $\int_0^{2\pi} \int_0^{\pi/2} \langle \cos \theta \sin^2 \varphi, \sin \theta \sin^2 \varphi, \sin \varphi \cos \varphi \rangle \cdot \langle \cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi \rangle d\varphi d\theta = \int_0^{2\pi} \int_0^{\pi/2} \sin \varphi d\varphi d\theta = \boxed{2\pi}$.
- (f) Parametrize by $\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, 1 - r^2 \rangle$ for $0 \leq \theta \leq 2\pi, 0 \leq r \leq 1$. Then $\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} = \langle 2r^2 \cos \theta, 2r^2 \sin \theta, r \rangle$ and $\mathbf{F} = \langle r^3 \sin^3 \theta, r^3 \cos^3 \theta, 1 \rangle$. Integral is then $\int_0^{2\pi} \int_0^1 \langle 2r^2 \cos \theta, 2r^2 \sin \theta, r \rangle \cdot \langle r^3 \sin^3 \theta, r^3 \cos^3 \theta, 1 \rangle dr d\theta = \int_0^{2\pi} \int_0^1 (2r^5 \sin \theta \cos \theta + r) dr d\theta = \boxed{\pi}$.
-
7. (a) $\nabla \cdot \mathbf{F} = \boxed{3x^2 + y + 3y^2 + x}$, $\nabla \times \mathbf{F} = \boxed{\langle 0, 0, y - x \rangle}$. Since $\nabla \times \mathbf{F} \neq \mathbf{0}$, field is not conservative.
- (b) $\nabla \cdot \mathbf{F} = \boxed{2}$, $\nabla \times \mathbf{F} = \boxed{\langle 0, 0, 0 \rangle}$. Since $\nabla \times \mathbf{F} = \mathbf{0}$, field is conservative. A potential is $U = \boxed{xyz + x^2 + 2yz}$.
- (c) $\nabla \cdot \mathbf{F} = \boxed{2xyz + 2x^2y}$, $\nabla \times \mathbf{F} = \boxed{\langle 0, x^2y - 4xyz, 2xz^2 - x^2z \rangle}$. Since $\nabla \times \mathbf{F} \neq \mathbf{0}$, field is not conservative.
- (d) $\nabla \cdot \mathbf{F} = \boxed{10yz + xe^z}$, $\nabla \times \mathbf{F} = \boxed{\langle 0, 0, 0 \rangle}$. Since $\nabla \times \mathbf{F} = \mathbf{0}$, field is conservative. A potential function is $U = \boxed{x^2yz + xe^z + 3y + 2z}$.
-
8. Integral is work done by $\mathbf{F} = \langle yz, xz, xy \rangle$ along C from $\mathbf{r}(0) = \langle 0, 0, 0 \rangle$ to $\mathbf{r}(1) = \langle e, \pi/4, \ln(2) \rangle$. \mathbf{F} is conservative with potential $U = xyz$, so work is simply $U(e, \pi/4, \ln(2)) - U(0, 0, 0) = \boxed{e \cdot (\pi/4) \cdot \ln(2)}$.
-
9. (a) We compute $\text{curl}(\mathbf{F}) = \langle 0, 0, 0 \rangle$. Since this is zero, \mathbf{F} is conservative.
- (b) Since \mathbf{F} is conservative and the particle travels around a closed path, the work is 0.